



Exclusion Guidelines

Condition	Exclude
Chickenpox	Until all blisters are crusted over (usually 5 to 6 days after rash started)
Cold Sores	No exclusion
Common Cold	No exclusion necessary if child feels well enough to participate in activities
CMV	No exclusion necessary
Diarrhea	Until diarrhea stops or until determined it is not a symptom of an infectious disease
Earache	No exclusion necessary
Fever	If temperature 100°F or higher (armpit) 2M a sore throat, rash, vomiting, diarrhea, pain, irritability, or confusion
Fifth Disease	No exclusion
Hand, Foot, and Mouth Disease	Only if the child is too sick to participate in activities (Coxsackie A)
Head Lice	Until nit free
Impetigo Rash	Until 24 hours after antibiotic treatment is started
Infectious Mononucleosis	No exclusion necessary if child feels well enough to participate in activities
Pinkeye Bacterial	Until 24 hours after antibiotic treatment is started
Viral	Until no drainage
Pinworms	Until treatment is started
Rashes	Until determined the rash is not a symptom of an infectious disease
Ringworm	Until treatment is started
Roseola	No exclusion
Rotavirus diarrhea	Until diarrhea stops
Scabies	Until treatment is started
Sore throat	Until determined it is not a symptom of an infectious disease
Strep throat	Until 24 hours after antibiotic treatment is started 2M fever is gone
Vomiting	Until determined the child is not infectious and not in danger of dehydration
Yeast Infections	Until treatment is started